



City Research Online

City, University of London Institutional Repository

Citation: Zhang, B., Shweikh, Y., Khawaja, A. P., Gallacher, J., Bauermeister, S., Foster, P. J., UKBiobank Eye and Vision Consortium, & Crabb, D. P. (2019). Associations with Corneal Hysteresis in a Population Cohort: Results from 96010 UK Biobank Participants. *Ophthalmology*, 126(11), pp. 1500-1510. doi: 10.1016/j.ophtha.2019.06.029

This is the accepted version of the paper.

This version of the publication may differ from the final published version.

Permanent repository link: <https://openaccess.city.ac.uk/id/eprint/26977/>

Link to published version: <https://doi.org/10.1016/j.ophtha.2019.06.029>

Copyright: City Research Online aims to make research outputs of City, University of London available to a wider audience. Copyright and Moral Rights remain with the author(s) and/or copyright holders. URLs from City Research Online may be freely distributed and linked to.

Reuse: Copies of full items can be used for personal research or study, educational, or not-for-profit purposes without prior permission or charge. Provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way.

Associations with corneal hysteresis in a population cohort: Results from 96,010 UK Biobank participants

Bing Zhang ¹, Yusrah Shweikh ², Anthony P Khawaja ², John Gallacher ¹, Sarah Bauermeister ¹, Paul J Foster ² on behalf of the UKBiobank Eye and Vision Consortium

Financial Support: None

Conflict of Interest: None

Running head: Associations with corneal hysteresis in UK Biobank

Abbreviations/Acronyms

CCT, central corneal thickness

CH, corneal hysteresis

CI, confidence interval

IOPg, Goldmann-correlated intraocular pressure

LOWESS, locally weighted scatterplot smoothing

OR, odds ratio

SLE, systemic lupus erythematosus

Author affiliations:

¹ Department of Psychiatry, University of Oxford, Oxford, UK

² NIHR Biomedical Research Centre, Moorfields Eye Hospital NHS Foundation Trust and UCL Institute of Ophthalmology, London, United Kingdom; Moorfields Eye Hospital, 162 City Road, London EC1V 2PD; UCL Institute of Ophthalmology, 11-43 Bath Street, London, EC1V 9EL

Corresponding author:

Prof Paul J Foster

Address: UCL Institute of Ophthalmology, 11-43 Bath Street, London, EC1V 9EL

Email: p.foster@ucl.ac.uk

Telephone: 07971 663189

Word count: 3152

FURTHER DETAILS

Authors' Contributions:

PJF, JG & BZ contributed to the conception and design of the study.

BZ performed data analysis.

All authors contributed to data interpretation.

All authors reviewed the results, read and critically revised the manuscript. All authors approved the final manuscript.

Declaration of interest (to be copied from ICMJE form once completed):

PJF reports personal fees from Allergan, Carl Zeiss, Google/DeepMind and Santen, a grant from Alcon, outside the submitted work;

APK, BZ, JG, SB, YS declare no competing interests.

Acknowledgements & Funding

Funding

The UK Biobank Eye and Vision Consortium is supported by grants from The NIHR Biomedical Research Centre at Moorfields Eye Hospital NHS Foundation Trust and UCL Institute of Ophthalmology, the Alcon Research Institute, Moorfields Eye Charity, and the International Glaucoma Association (UK).

No funders had a direct role in the collection, management, analysis, or interpretation of the data; preparation, review, or approval of the manuscript; nor in the decision to submit the manuscript for publication.

Acknowledgements

PJF received support from the Richard Desmond Charitable Trust, via Fight for Sight, London. APK is supported by a Moorfields Eye Charity Career Development Fellowship. PJF & APK received salary support from the NIHR BRC at Moorfields Eye Hospital. These authors acknowledge a proportion of our financial support from the UK Department of Health through an award made by the National Institute for Health Research to Moorfields Eye Hospital NHS Foundation Trust and UCL Institute of Ophthalmology for a Biomedical Research Centre for Ophthalmology.

Ethical approval: The North West Multi-centre Research Ethics Committee approved the study (reference no., 06/MRE08/65), in accordance with the tenets of the Declaration of Helsinki. Detailed information about the study is available at the UK Biobank web site (www.ukbiobank.ac.uk)

1 **Abstract**

2 **Purpose:** To describe the distribution of corneal hysteresis (CH) in a large cohort and explore its
3 associated factors and possible clinical applications.

4 **Design:** Cross-sectional study within the UK Biobank, a large cohort study in the United Kingdom.

5 **Participants:** We analyzed CH data from 93,345 eligible participants in the UK Biobank cohort,
6 aged 40 to 69 years.

7 **Methods:** All analyses were performed using left eye data. Linear regression models were used to
8 evaluate associations between CH and demographic, lifestyle, ocular and systemic variables.
9 Piecewise logistic regression models were used to explore the relationship between self-reported
10 glaucoma and CH.

11 **Main outcome measures:** CH (mmHg).

12 **Results:** The mean CH was 10.6 mmHg (10.4 mmHg in males and 10.8 mmHg in females). After
13 adjusting for covariables, CH was significantly negatively associated with male sex, age, Black
14 ethnicity, self-reported glaucoma, diastolic blood pressure and height. CH was significantly
15 positively associated with smoking, hyperopia, diabetes, systemic lupus erythematosus (SLE),
16 greater deprivation (Townsend index) and Goldmann-correlated intraocular pressure (IOPg). Self-
17 reported glaucoma and CH were significantly associated when CH was less than 10.1mmHg (OR
18 0.86, 95%CI 0.79-0.94 per mmHg CH increase) after adjusting for covariables. When CH exceeded
19 10.1 mmHg, there was no significant association between CH and self-reported glaucoma.

20 **Conclusion:** In our analyses, CH was significantly associated with factors including age, sex and
21 ethnicity which should be taken into account when interpreting CH values. In our cohort, lower CH
22 was significantly associated with a higher prevalence of self-reported glaucoma when CH was less

23 than 10.1mmHg. CH may serve as a biomarker aiding glaucoma case detection.

It is well recognized that variation in central corneal thickness (CCT) influences the accuracy of intraocular pressure (IOP) measurements¹⁻³. It has also been hypothesized that CCT independently influences the risk of glaucoma, with thin CCT evidenced in those at highest risk⁴. However, this view is not universally accepted, as one particular high-risk group (African Americans) typically have thinner CCT than people of European heritage⁵. A plausible alternative explanation is that thin CCT is a biomarker for race, and identifies those at highest risk, attributable to other ocular or systemic factors.

Corneal hysteresis (CH) offers an alternative index of corneal biomechanical characteristics to CCT and reflects the viscoelastic damping effect of corneal tissues, defined as the difference in air pulse pressure between inward and outward applanation forces^{6,7}. Recent evidence indicates CH can also provide valuable information related to the presence, progression and response to therapy of glaucoma^{8,9}. CH can be measured simultaneously with IOP using non-contact tonometry with augmented functionality. Differences in CH have been reported not only in glaucoma but also in many systemic diseases including thyroid eye disease¹⁰, rheumatoid arthritis¹¹, psoriasis¹², acromegaly¹³ and myotonic dystrophy¹⁴, which suggests CH may play a clinical role in fields other than ophthalmology. Previous studies on CH are limited by small sample sizes^{15,16}. The distribution of CH and its associations with demographic, ocular and systemic variables remain to be accurately determined and confirmed in a large sample.

The UK Biobank is one of the largest prospective population cohort studies in the world. In this study, we aimed to report the distribution of CH by age, sex and ethnicity, and explore its associations including the relationship between CH and self-reported glaucoma. We also tested the association between CH and 16 self-reported diseases selected based on existing literature¹⁰⁻¹³.

Methods

Study population

The UK Biobank is a multisite community-based cohort study with 502,544 participants. All UK residents aged 40 to 69 who registered with the National Health Service and lived within 25 miles of any of the 22 assessment centers were invited to join the study. The initial visit assessments took place between 2006 and 2010. Eye assessments were carried out from 2009 in 6 recruitment centers (5 in England and 1 in Wales) which enrolled 133,953 participants. The UK Biobank study was approved by the North West Multi-centre Research Ethics Committee (Reference No. 06/MRE08/65) and adhered to the tenets of the Declaration of Helsinki. Written consent was obtained from every participant. More detailed information and protocols for UK Biobank are available online (<http://www.ukbiobank.ac.uk/>).

Ethnicity was self-reported by participants and selected from White, Asian, Black, Chinese, mixed and other ethnic backgrounds. Socioeconomic status was derived using the Townsend deprivation index estimated using residence postcodes. This represents an indicative measure of economic deprivation in an area and higher scores indicate worse socioeconomic status¹⁷.

Measurements

Cohort characteristics and ophthalmic measures have been previously described¹⁸. Visual acuity was measured using a bespoke computerized logMAR acuity measure conforming to British Standard BS4274-1968¹⁹, with left eye following right eye. Autorefractometry was performed with the RC5000 Auto Refractometer (Tomey, Japan). After measuring visual acuity and refraction, CH and Goldmann-correlated IOP (IOPg) were measured with the Reichert Ocular Response Analyser (ORA, Reichert, Inc. USA) according to a predetermined protocol (available online

<http://biobank.ctsu.ox.ac.uk/crystal/refer.cgi?id=100236>). Participants who had any eye surgery within the preceding 4 weeks were excluded from tests. The measurements were performed first in the right eye and taken only once in each eye. If participants blinked during the test a further measurement was attempted.

Blood pressure was measured with an automatic blood pressure monitor, HEM-70151T (Omron, The Netherlands). Two measurements were performed for each participant and the average was used for analysis if the values of both were available. Height was measured with the Seca 202 instrument (Seca, UK).

Medical History

All diseases were self-reported by participants via verbal interviews conducted by trained nurses or via touchscreen questionnaires. Self-reported eye disorder(s) status was collected in the verbal interview or was selected by participants from a list of eye disorders in response to the question “Has a doctor told you that you have any of the following problems with your eyes?”. The list of eye disorders was:

1. Diabetes related eye disease
2. Glaucoma
3. Injury or trauma resulting in loss of vision
4. Cataract
5. Macular degeneration
6. Other serious eye condition
7. None of the above
8. Prefer not to answer

9. Do not know

Smoking and alcohol consumption were self-reported via touchscreen questionnaires. Smoking status was trichotomized for the purpose of analysis to current smokers, ex-smokers and those that have never smoked. Alcohol consumption was pentachotomized to daily/almost daily, weekly or more often, monthly or more often, occasional and never. The use of IOP lowering medications was recorded by trained interviewers. Only currently and regularly used ones were recorded. IOP lowering medication status was dichotomized to user and non-user for analysis.

More detailed information about all variables is available online

(<http://biobank.ctsu.ox.ac.uk/crystal/index.cgi>).

Eligibility criteria

All participants who had available ORA data (CH and IOPg) in the left eye were used for this analysis. Participants who met any exclusion criteria in Figure 1 were excluded from the analyses. 0.5% of participants who were younger than 40 or older than 69 years were excluded based on the UK Biobank eligibility criteria. Extreme values (lowest 0.5% and highest 0.5%) of CH and IOPg may represent measurement errors and were therefore excluded. We excluded participants with a history of eye injury in their left eye, diabetes related eye disease, macular degeneration or other serious eye conditions (except for glaucoma and cataract) in either eye. Left eyes without data on ocular comorbidities and/or refractive error, and/or with high refractive errors (spherical equivalent $>+5D$ or $<-6D$) and/or high astigmatism (absolute value of cylindrical power $>3D$) and/or a history of refractive surgery were excluded. Participants with a history of surgery or laser for glaucoma or ocular hypertension were also excluded. Of the 93,345 left eyes remained in analysis, 1,208 eyes with self-reported glaucoma were excluded for analyses of CH distribution.

Statistical analysis

All analyses were performed using left eye data which were captured after right eye data as specified in the study protocol. This may mean left eye data are less prone to artefact, such as blinking, in our cohort²⁰. We included refractive error in analyses as the spherical equivalent in dioptres (D, sphere power+1/2 cylinder power). For glaucoma status, controls were defined as participants without self-reported glaucoma in either eye.

A descriptive analysis of CH in left eyes stratified by age, sex and ethnicity was conducted after excluding all participants with self-reported glaucoma. One-way analysis of variance was performed to compare means of CH by age, sex and ethnicity.

Associations between CH and other demographic, ocular and systemic factors and self-reported glaucoma were evaluated with univariable linear regression and all factors with $p<0.05$ in univariable analysis were also analyzed with multivariable linear regression.

We analyzed the relationship between self-reported glaucoma and CH using the following steps:

- 1) Locally weighted scatterplot smoothing (LOWESS)²¹, a method usually used to visualize the structure of data²², was used to explore the relationship between self-reported glaucoma and corneal hysteresis. The turning point(s) found on the LOWESS curve was used as node(s) for piecewise analysis.
- 2) Piecewise logistic regression for self-reported glaucoma and CH was performed in three models after adjusting for covariables.
- 3) The joint distribution of the proportion of self-reported glaucoma, CH and IOPg was displayed using a 3D bar chart.

We then applied linear regression to evaluate the relationships between CH and 16 systemic diseases

after adjusting for covariables.

The 3D bar chart was plotted using Excel for Office 365 (MicrosoftCorp, CA, USA). All other analyses were performed and plots generated using STATA/SE-15 (StataCorp LLC, TX, USA).

Results

All analyses were performed using left eye data in this study. 111,942 UK Biobank participants had available CH values for left eyes. After data cleaning as shown in Figure 1, the mean CH was 10.60 ± 1.88 mmHg (95% CI 10.59-10.62 mmHg) in the 92,137 eyes without self-reported glaucoma.

The distribution of mean CH stratified by age, sex and ethnicity is summarized in Table 1. A significant difference in CH was found between participants with different ethnicities ($p < 0.001$).

CH values were lower in Black people (9.62 ± 1.87 mmHg, 95% CI 9.56-9.69 mmHg) compared to White participants (10.66 ± 1.87 mmHg, 95% CI 10.65-10.67 mmHg). CH was significantly greater in females (10.79 ± 1.86 mmHg, 95% CI 10.77-10.80 mmHg) compared to males (10.39 ± 1.88 mmHg, 95% CI 10.37-10.40 mmHg, $p < 0.001$). Overall, CH was also significantly higher in younger people across the whole age spectrum enrolled (mean 10.91 ± 1.91 mmHg, 95% CI 10.87-10.95 mmHg for those aged 40-44 compared to 10.30 ± 1.84 mmHg, 95% CI 10.27-10.32 mmHg for those aged 65-69, $p < 0.001$).

The associations of CH were analyzed with linear regression models as shown in Table 2. CH was significantly associated with all included factors except for visual acuity and alcohol intake frequency. In the multivariable linear regression model after adjusting for covariates, CH was significantly higher in women (0.193 mmHg, $p = 2.07 \times 10^{-27}$), smokers (reference: never smoked; 0.095 mmHg former smokers, $p = 7.71 \times 10^{-13}$; 0.419 mmHg current smokers, $p = 1.22 \times 10^{-84}$), participants with a higher Townsend deprivation index (0.012 mmHg/Unit, $p = 7.82 \times 10^{-8}$) and self-

reported diabetes (0.283 mmHg , $p=1.25 \times 10^{-20}$). CH was significantly lower in older participants (-0.033 mmHg/year , $p=0$), Black participants (reference: white; -1.219 mmHg , $p=1.03 \times 10^{-260}$), Asian participants (reference: white; -0.461 mmHg , $p=2.08 \times 10^{-45}$), participants with higher blood pressure ($-0.0076 \text{ mmHg/1mmHg diastolic blood pressure}$, $p=1.29 \times 10^{-33}$), greater height (-0.016 mmHg/cm , $p=4.71 \times 10^{-61}$), greater myopia (0.034 mmHg/D , $p=3.06 \times 10^{-26}$) and in those with self-reported glaucoma (-0.516 mmHg , $p=1.13 \times 10^{-15}$).

Figure 2, Table 3 and Figure 3 show the relationship between self-reported glaucoma and CH. Overall, lower CH was associated with a higher proportion of self-reported glaucoma. As shown in Figure 2A, when CH was less than approximately 10 mmHg , the proportion of self-reported glaucoma increased markedly when CH decreased. However, with increases in CH above 10 mmHg the proportion of self-reported glaucoma remained relatively stable at around 1%. The LOWESS curve shapes were similar in analyses stratified by age (Figure 2B) and IOPg (Figure 2C), with sharp rises in the proportions of self-reported glaucoma at CH values less than approximately 10 mmHg . Piecewise logistic regressions were performed with a node set at 10.1 mmHg (Table 3). As shown in the online supplementary material, 10.1 mmHg was the smallest node that self-reported glaucoma and CH were significantly associated when CH was less than the node while there was no association between self-reported glaucoma and CH when CH was greater than the 10.1 mmHg node in all three models. When CH was less than 10.1 mmHg , higher CH was a protective factor for self-reported glaucoma. A 1 mmHg increase in CH was associated with an OR of 0.78 (95% CI $0.73\text{-}0.82$, $p<0.001$) after adjusting for age, sex and ethnicity in Model I, an OR of 0.82 (95% CI $0.78\text{-}0.87$, $p<0.001$) in Model II (Model I with further adjusting for IOPg) and an OR of 0.86 (95% CI $0.79\text{-}0.94$, $p<0.001$) in Model III (the maximally adjusted model). When CH exceeded 10.1

mmHg it was not associated with self-reported glaucoma in all three models (Table 3).

The relationship between self-reported glaucoma, CH and IOPg is displayed using a 3D bar chart (Figure 3). In keeping with the analyses reported in Figure 2C and Table 3, the proportion of self-reported glaucoma was highest in participants with high IOPg and low CH, and lowest in the participants whose IOPg was not high and CH was not low.

We analyzed associations between CH and 16 self-reported disorders of the thyroid gland, pituitary gland and other immunological/systemic disorders (Table 4). Only systemic lupus erythematosus (SLE) was significantly associated with CH following correction for multiple testing ($p < 0.003125$, Bonferroni-corrected threshold). CH was significantly higher in participants with self-reported SLE (0.549, 95% CI 0.237-0.862 mmHg in the fully adjusted model).

Discussion

In this large UK cohort, we have described mean CH stratified by age, sex and ethnicity (Table 1).

We found that CH was significantly lower in Black participants and in older age groups, which is consistent with previously published findings^{15,23}. Past studies indicate that CH and CCT are positively associated²⁴⁻²⁶ and CCT is negatively associated with darker skin pigmentation²⁷. One explanation for the variation in CH by ethnicity may be differences mediated by changes in CCT. Conversely, previous publications revealed no significant association between CCT and age^{7,28,29}, suggesting an independent association between lower CH and older age.

CH was significantly higher in smokers in our cohort (both current and former smokers). A previous, smaller study had suggested this but results were inconclusive³⁰. The mechanisms underlying the relationship between smoking and corneal changes are unknown^{31,32} and the association between smoking and corneal ectatic disorders is controversial^{33,34}. An epidemiological study showed a

marked reduction in the incidence of keratoconus amongst smokers³⁴, implying altered corneal biomechanics. This is supported by experimental evidence of collagen crosslinking by formaldehyde, a constituent of cigarette smoke, with resulting increased resistance to collagenases³⁴. Smoking has also been reported to damage the tear film^{35,36} and possibly the corneal endothelium³⁷, which may influence CCT and CH measurements. We found no significant association between alcohol consumption and CH.

Our findings in Figure 2, Table 3 and Figure 3 suggest that CH may be useful in glaucoma risk stratification in clinical practice. Figure 2 and Table 3 indicate that a CH value of 10.1 mmHg could play a role as cutoff point in clinical practice to evaluate a patient's risk of glaucoma. When CH is less than 10.1mmHg, lower CH may be associated with a higher risk of glaucoma (OR 1.16, 95% CI 1.07-1.26 per mmHg CH decrease in the fully adjusted model). When CH was greater than 10.1mmHg, the rate of self-reported glaucoma remained relatively stable with further increases in CH. Medeiros et al reported that lower CH with values below 10mmHg was a risk factor for glaucoma progression³⁸.

CH measurement demonstrates good repeatability³⁹ and there are no significant diurnal fluctuations^{26,40}, making CH measurement a potentially attractive addition to current glaucoma risk stratification methods. CH has been shown to be lower in different types of glaucoma including open angle glaucoma, angle closure glaucoma, normal tension glaucoma, pseudoexfoliative glaucoma and congenital glaucoma⁴¹⁻⁴⁶. Lower CH is also positively associated with visual field progression^{8,38}.

Some studies have found a positive association between CH and glaucoma-related changes in optic disc morphology⁴⁷⁻⁴⁹ whereas others found no such relationship⁵⁰⁻⁵². Unlike CH, IOP and CCT measurements are limited by significant diurnal variation^{26,40,53-55}. Figure 2C, Table 3 and Figure 3

show that CH and IOPg could be analyzed together in clinical settings to evaluate glaucoma risk, as the risk of self-reported glaucoma was highest in participants with low CH and high IOPg, and lowest in participants whose IOPg was not high and CH was not low.

In analyses for associations between CH and self-reported disorders shown in Table 4, only SLE was significantly associated with CH at $p<0.003$ (Bonferroni-corrected threshold for multiple testing). We found that CH was significantly higher in participants with SLE, which is contradictory to the result in a case-control study which reported CH was lower in SLE patients⁵⁶. Lower CH has also been reported in thyroid eye disease¹⁰, however we did not find an association between CH and thyroid disorders. We also did not find associations between CH and rheumatoid arthritis or psoriasis as previously published^{11,12}. Participants with acromegaly in our cohort had higher CH values (at $p<0.05$), in agreement with findings from Ozkok and colleagues¹³, however our result was not significant after correction for multiple testing. Our study also shows higher CH amongst patients with diabetes as previously reported^{57,58}. Former studies have yielded variable results when evaluating CH in diabetes⁵⁸⁻⁶¹.

The very large sample size and standardized techniques are major strengths of our study, allowing us to detect and quantify small effects. However, the study is limited by the fact that all disease statuses were self-reported by participants which can result in misclassification error⁶². UK Biobank has a low response rate of 5.5% which limits external validity. With respect to glaucoma, there will be an under-ascertainment of disease since approximately 50% of cases may not have been diagnosed⁶². Meanwhile participants with ocular hypertension, suspected glaucoma or cataracts may report a diagnosis of glaucoma. The potential impact of these errors is unknown. We excluded participants with a past history of surgery or laser for glaucoma or ocular hypertension. A potential

confounding variable in the reported association between CH and glaucoma is the use of IOP lowering medications, which may significantly alter corneal biomechanical properties^{9,63,64}. The binary variable of current, regular IOP lowering medication use versus no use in this study may oversimplify the effects of different medications on corneal biomechanics. CH and IOPg in this study were measured together using the same instrument and adjusting one for the other makes interpretation difficult. Despite this, we found weak correlation between them ($\rho=0.045$) in the sample after data cleaning. Investigation into the association between CH and diseases including glaucoma, SLE and diabetes is scarce and we anticipate that future research will build on our findings.

Our study offers CH reference values for future research and clinical practice. We also report associations between CH and age, sex, ethnicity, smoking status, refractive error, self-reported glaucoma, diabetes and SLE, which may be important when interpreting CH. CH measurement may play a role in clinical practice for glaucoma and other ocular and systemic conditions.

References

1. Ehlers N, Bramsen T, Sperling S. Applanation tonometry and central corneal thickness. *Acta Ophthalmol.* 1975;53(1):34-43.
2. Whitacre MM, Stein R. Sources of error with use of Goldmann-type tonometers. *Surv Ophthalmol.* 1993;38(1):1-30.
3. Whitacre MM, Stein RA, Hassanein K. The effect of corneal thickness on applanation tonometry. *Am J Ophthalmol.* 1993;115(5):592-596.
4. Gordon MO, Beiser JA, Brandt JD, et al. The Ocular Hypertension Treatment Study: baseline factors that predict the onset of primary open-angle glaucoma. *Arch Ophthalmol.* 2002;120(6):714-720.
5. Brandt JD, Beiser JA, Kass MA, Gordon MO. Central corneal thickness in the Ocular Hypertension Treatment Study (OHTS). *Ophthalmology.* 2001;108(10):1779-1788.
6. Luce DA. Determining in vivo biomechanical properties of the cornea with an ocular response analyzer. *J Cataract Refract Surg.* 2005;31(1):156-162.
7. Hoffmann EM, Lamparter J, Mirshahi A, et al. Distribution of central corneal thickness and its

273 association with ocular parameters in a large central European cohort: the Gutenberg health
274 study. *PloS One*. 2013;8(8):e66158.

275 8. Congdon NG, Broman AT, Bandeenroche K, Grover D, Quigley HA. Central corneal thickness
276 and corneal hysteresis associated with glaucoma damage. *Am J Ophthalmol*. 2006;141(5):868-
277 875.

278 9. Agarwal DR, Ehrlich JR, Shimmyo M, Radcliffe NM. The relationship between corneal
279 hysteresis and the magnitude of intraocular pressure reduction with topical prostaglandin
280 therapy. *Br J Ophthalmol*. 2012;96(2):254-257.

281 10. Karabulut GO, Kaynak P, Altan C, et al. Corneal biomechanical properties in thyroid eye disease.
282 *Kaohsiung J Med Sci*. 2014;30(6):299-304.

283 11. Can ME, Erten S, Can GD, Cakmak HB, Sarac O, Cagil N. Corneal biomechanical properties
284 in rheumatoid arthritis. *Eye Contact Lens*. 2015;41(6):382-385.

285 12. Celik U, Aykut V, Celik B, et al. A comparison of corneal biomechanical properties in patients
286 with psoriasis and healthy subjects. *Eye Contact Lens*. 2015;41(2):127-129.

287 13. Ozkok A, Hatipoglu E, Tamcelik N, et al. Corneal biomechanical properties of patients with
288 acromegaly. *Br J Ophthalmol*. 2014;98(5):651-657.

289 14. Garcia Filho CA, Prata TS, Sousa AK, Doi LM, Melo Jr LA. Intraocular pressure, corneal
290 thickness, and corneal hysteresis in Steinert's myotonic dystrophy. *Arq Bras Oftalmol*.
291 2011;74(3):161-162.

292 15. Haseltine SJ, Pae J, Ehrlich JR, Shammam M, Radcliffe NM. Variation in corneal hysteresis and
293 central corneal thickness among black, hispanic and white subjects. *Acta Ophthalmol*.
294 2012;90(8):e626-e631.

295 16. Wang JK, Huang TL, Pei-Yuan S, Chang PY. Factors affecting corneal hysteresis in Taiwanese
296 adults. *Eye Sci*. 2015;30(3):89-93.

297 17. Rice LJ, Jiang C, Wilson SM, Burwell-Naney K, Samantapudi A, Zhang H. Use of segregation
298 indices, Townsend Index, and air toxics data to assess lifetime cancer risk disparities in
299 metropolitan Charleston, South Carolina, USA. *Int J Environ Res Public Health*.
300 2014;11(5):5510-5526.

301 18. Chua SYL, Thomas D, Allen N, et al. Cohort profile: design and methods in the eye and vision
302 consortium of UK Biobank. *BMJ open*. 2019;9(2):e025077.

303 19. Standard B. Test charts for determining distance visual acuity: BS 4274-1968. *London: British*
304 *Standards Institution*. 1968.

305 20. Chan MP, Grossi CM, Khawaja AP, et al. Associations with intraocular pressure in a large cohort:
306 results from the UK Biobank. *Ophthalmology*. 2016;123(4):771-782.

307 21. Cleveland WS. Robust locally weighted regression and smoothing scatterplots. *J Am Stat Assoc*.
308 1979;74(368):829-836.

309 22. Abu-Hanna A, de Keizer N. Integrating classification trees with local logistic regression in
310 Intensive Care prognosis. *Artif Intell Med*. 2003;29(1-2):5-23.

311 23. Celebi ARC, Kilavuzoglu AE, Altiparmak UE, Cosar Yurteri CB. Age-related change in corneal
312 biomechanical parameters in a healthy Caucasian population. *Ophthalmic Epidemiol*.
313 2018;25(1):55-62.

314 24. Shah S, Laiquzzaman M, Cunliffe I, Mantry S. The use of the Reichert ocular response analyser
315 to establish the relationship between ocular hysteresis, corneal resistance factor and central
316 corneal thickness in normal eyes. *Contact Lens Anterio*. 2006;29(5):257-262.

- 317 25. Mangouritsas G, Morphis G, Mourtzoukos S, Feretis E. Association between corneal hysteresis
318 and central corneal thickness in glaucomatous and non-glaucomatous eyes. *Acta Ophthalmol.*
319 2009;87(8):901-905.
- 320 26. Kida T, Liu JHK, Weinreb RN. Effects of aging on corneal biomechanical properties and their
321 impact on 24-hour measurement of intraocular pressure. *Am J Ophthalmol.* 2008;146(4):567-
322 572.
- 323 27. Dimasi DP, Hewitt AW, Kagame K, et al. Ethnic and mouse strain differences in central corneal
324 thickness and association with pigmentation phenotype. *PloS One.* 2011;6(8):e22103.
- 325 28. Zheng Y, Huang G, Huang W, He M. Distribution of central and peripheral corneal thickness in
326 Chinese children and adults: the Guangzhou twin eye study. *Cornea.* 2008;27(7):776-781.
- 327 29. Wolfs RC, Klaver CC, Vingerling JR, Grobbee DE, Hofman A, de Jong PT. Distribution of
328 central corneal thickness and its association with intraocular pressure: The Rotterdam Study.
329 *Am J Ophthalmol.* 1997;123(6):767-772.
- 330 30. Kilavuzoglu AE, Celebi AR, Altiparmak UE, Cosar CB. The effect of smoking on corneal
331 biomechanics. *Curr Eye Res.* 2017;42(1):16-20.
- 332 31. Madhukumar E, Vijayammal PL. Influence of cigarette smoke on cross-linking of dermal
333 collagen. *Indian J Exp Biol.* 1997;35(5):483-486.
- 334 32. Wollensak G, Spoerl E. Collagen crosslinking of human and porcine sclera. *J Cataract Refract*
335 *Surg.* 2004;30(3):689-695.
- 336 33. Jonas JB, Nangia V, Matin A, Kulkarni M, Bhojwani K. Prevalence and associations of
337 keratoconus in rural maharashtra in central India: the central India eye and medical study. *Am J*
338 *Ophthalmol.* 2009;148(5):760-765.
- 339 34. Raiskup-Wolf F, Spoerl E, Kuhlisch E, Pillunat LE. Cigarette smoking is negatively associated
340 with keratoconus. *J Refract Surg.* 2008;24(7):S737-S740.
- 341 35. Altinors DD, Akça S, Akova YA, et al. Smoking associated with damage to the lipid layer of the
342 ocular surface. *Am J Ophthalmol.* 2006;141(6):1016-1021.
- 343 36. Yoon K-C, Song B-Y, Seo M-S. Effects of smoking on tear film and ocular surface. *Korean*
344 *Journal of Ophthalmology.* 2005;19(1):18-22.
- 345 37. Sayin N, Kara N, Pekel G, Altinkaynak H. Effects of chronic smoking on central corneal
346 thickness, endothelial cell, and dry eye parameters. *Cutan Ocul Toxicol.* 2014;33(3):201-205.
- 347 38. Medeiros FA, Freitas D, Lisboa R, Kuang TM, Zangwill LM, Weinreb RN. Corneal hysteresis
348 as a risk factor for glaucoma progression: a prospective longitudinal study. *Ophthalmology.*
349 2013;120(8):1533-1540.
- 350 39. David VP, Stead RE, Vernon SA. Repeatability of ocular response analyzer metrics: a gender-
351 based study. *Optom Vis Sci.* 2013;90(7):691-699.
- 352 40. Kotecha A, Crabb DP, Spratt A, Garway-Heath DF. The relationship between diurnal variations
353 in intraocular pressure measurements and central corneal thickness and corneal hysteresis.
354 *Invest Ophthalmol Vis Sci.* 2009;50(9):4229-4236.
- 355 41. Abitbol O, Bouden J, Doan S, Hoang-Xuan T, Gatinel D. Corneal hysteresis measured with the
356 Ocular Response Analyzer in normal and glaucomatous eyes. *Acta Ophthalmol.*
357 2010;88(1):116-119.
- 358 42. Castro DP, Prata TS, Lima VC, Biteli LG, de Moraes CG, Jr PA. Corneal viscoelasticity
359 differences between diabetic and nondiabetic glaucomatous patients. *J Glaucoma.*
360 2010;19(5):341-343.

- 361 43. Kaushik S, Pandav SS, Banger A, Aggarwal K, Gupta A. Relationship between corneal
362 biomechanical properties, central corneal thickness, and intraocular pressure across the
363 spectrum of glaucoma. *Am J Ophthalmol*. 2012;153(5):840-849.
- 364 44. Narayanaswamy A, Su DH, Baskaran M, et al. Comparison of ocular response analyzer
365 parameters in Chinese subjects with primary angle-closure and primary open-angle glaucoma.
366 *Arch Ophthalmol*. 2011;129(4):429-434.
- 367 45. Ozkok A, Tamcelik N, Ozdamar A, Sarici AM, Cicik E. Corneal viscoelastic differences
368 between pseudoexfoliative glaucoma and primary open-angle glaucoma. *J Glaucoma*.
369 2013;22(9):740-745.
- 370 46. Gatzoufas Z, Labiris G, Stachs O, et al. Biomechanical profile of the cornea in primary
371 congenital glaucoma. *Acta Ophthalmol*. 2013;91(1):e29-e34.
- 372 47. Bochmann F, Ang GS, Azuara-Blanco A. Lower corneal hysteresis in glaucoma patients with
373 acquired pit of the optic nerve (APON). *Graefes Arch Clin Exp Ophthalmol*. 2008;246(5):735.
- 374 48. Khawaja AP, Chan MP, Broadway DC, et al. Corneal biomechanical properties and glaucoma-
375 related quantitative traits in the EPIC-Norfolk Eye Study. *Invest Ophthalmol Vis Sci*.
376 2014;55(1):117-124.
- 377 49. Prata TS, Lima VC, Guedes LM, et al. Association between corneal biomechanical properties
378 and optic nerve head morphology in newly diagnosed glaucoma patients. *Clin Exp Ophthalmol*.
379 2012;40(7):682-688.
- 380 50. Carbonaro F, Hysi PG, Fahy SJ, Nag A, Hammond CJ. Optic disc planimetry, corneal hysteresis,
381 central corneal thickness, and intraocular pressure as risk factors for glaucoma. *Am J*
382 *Ophthalmol*. 2014;157(2):441-446.
- 383 51. Mansouri K, Leite MT, Weinreb RN, Tafreshi A, Zangwill LM, Medeiros FA. Association
384 between corneal biomechanical properties and glaucoma severity. *Am J Ophthalmol*.
385 2012;153(3):419.
- 386 52. Vu DM, Silva FQ, Haseltine SJ, Ehrlich JR, Radcliffe NM. Relationship between corneal
387 hysteresis and optic nerve parameters measured with spectral domain optical coherence
388 tomography. *Graefes Arch Clin Exp Ophthalmol*. 2013;251(7):1777-1783.
- 389 53. Harper CL, Boulton ME, Bennett D, et al. Diurnal variations in human corneal thickness. *Br J*
390 *Ophthalmol*. 1996;80(12):1068-1072.
- 391 54. du Toit R, Vega JA, Fonn D, Simpson T. Diurnal variation of corneal sensitivity and thickness.
392 *Cornea*. 2003;22(3):205-209.
- 393 55. Sharifipour F, Farrahi F, Moghaddasi A, Idani A, Yaseri M. Diurnal variations in intraocular
394 pressure, central corneal thickness, and macular and retinal nerve fiber layer thickness in
395 diabetics and normal individuals. *J Ophthalmic Vis Res*. 2016;11(1):42-47.
- 396 56. Yazici AT, Kara N, Yüksel K, et al. The biomechanical properties of the cornea in patients with
397 systemic lupus erythematosus. *Eye*. 2011;25(8):1005-1009.
- 398 57. Hussnain SA, Alsberge JB, Ehrlich JR, Shimmyo M, Radcliffe NM. Change in corneal
399 hysteresis over time in normal, glaucomatous and diabetic eyes. *Acta Ophthalmol*.
400 2015;93(8):e627-e630.
- 401 58. Scheler A, Spoerl E, Boehm AG. Effect of diabetes mellitus on corneal biomechanics and
402 measurement of intraocular pressure. *Acta Ophthalmol*. 2012;90(6):e447-e451.
- 403 59. Goldich Y, Barkana Y, Gerber Y, et al. Effect of diabetes mellitus on biomechanical parameters
404 of the cornea. *J Cataract Refract Surg*. 2009;35(4):715-719.

- 405 60. Kotecha A, Oddone F, Sinapis C, et al. Corneal biomechanical characteristics in patients with
406 diabetes mellitus. *J Cataract Refr Surg*. 2010;36(11):1822-1828.
- 407 61. Sahin A, Bayer A, Ozge G, Mumcuoglu T. Corneal biomechanical changes in diabetes mellitus
408 and their influence on intraocular pressure measurements. *Invest Ophthalmol Vis Sci*.
409 2009;50(10):4597-4604.
- 410 62. Shweikh Y, Ko F, Chan MP, et al. Measures of socioeconomic status and self-reported glaucoma
411 in the U.K. Biobank cohort. *Eye*. 2015;29(10):1360-1367.
- 412 63. Sun L, Shen M, Wang J, et al. Recovery of corneal hysteresis after reduction of intraocular
413 pressure in chronic primary angle-closure glaucoma. *Am J Ophthalmol*. 2009;147(6):1061-1066.
- 414 64. Tsikripis P, Papaconstantinou D, Koutsandrea C, Apostolopoulos M, Georgalas I. The effect of
415 prostaglandin analogs on the biomechanical properties and central thickness of the cornea of
416 patients with open-angle glaucoma: a 3-year study on 108 eyes. *Drug Des Devel Ther*.
417 2013;7:1149-1156.

418

419 **UK Biobank Eye and Vision Consortium membership**

420 UK Biobank Eye & Vision Consortium: The UK Biobank Eye & Vision Consortium members are Tariq
421 Aslam, PhD, Manchester University, Sarah A. Barman, PhD, Kingston University, Jenny H. Barrett, PhD,
422 University of Leeds, Paul Bishop, PhD, Manchester University, Peter Blows, BSc, NIHR Biomedical
423 Research Centre, Catey Bunce, DSc, King's College London, Roxana O. Carare, PhD, University of
424 Southampton, Usha Chakravarthy, FRCOphth, Queens University Belfast, Michelle Chan, FRCOphth,
425 NIHR Biomedical Research Centre, Sharon Y.L. Chua, PhD, NIHR Biomedical Research Centre, David
426 P. Crabb, PhD, UCL, Philippa M. Cumberland, MSc, UCL Great Ormond Street Institute of Child Health,
427 Alexander Day, PhD, NIHR Biomedical Research Centre, Parul Desai, PhD, NIHR Biomedical Research
428 Centre, Bal Dhillon, FRCOphth, University of Edinburgh, Andrew D. Dick, FRCOphth, University of
429 Bristol, Cathy Egan, FRCOphth, NIHR Biomedical Research Centre, Sarah Ennis, PhD, University of
430 Southampton, Paul Foster, PhD, NIHR Biomedical Research Centre, Marcus Fruttiger, PhD, NIHR
431 Biomedical Research Centre, John E.J. Gallacher, PhD, University of Oxford, David F. GARWAY-
432 HEATH FRCOphth- NIHR Biomedical Research Centre, Jane Gibson, PhD, University of Southampton,
433 Dan Gore, FRCOphth, NIHR Biomedical Research Centre, Jeremy A. Guggenheim, PhD, Cardiff

434 University, Chris J. Hammond, FRCOphth, King's College London, Alison Hardcastle, PhD, NIHR
435 Biomedical Research Centre, Simon P. Harding, MD, University of Liverpool, Ruth E. Hogg, PhD,
436 Queens University Belfast, Pirro Hysi, PhD, King's College London, Pearse A. Keane, MD, NIHR
437 Biomedical Research Centre, Sir Peng T. Khaw, PhD, NIHR Biomedical Research Centre, Anthony P.
438 Khawaja, DPhil, NIHR Biomedical Research Centre, Gerassimos Lascaratos, PhD, NIHR Biomedical
439 Research Centre, Andrew J. Lotery, MD, University of Southampton, Tom Macgillivray, PhD, University
440 of Edinburgh, Sarah Mackie, PhD, University of Leeds, Keith Martin, FRCOphth, University of
441 Cambridge, Michelle McGaughey, Queen's University Belfast, Bernadette McGuinness, PhD, Queen's
442 University Belfast, Gareth J. McKay, PhD, Queen's University Belfast, Martin McKibbin, FRCOphth,
443 Leeds Teaching Hospitals NHS Trust, Danny Mitry, PhD, NIHR Biomedical Research Centre, Tony
444 Moore, FRCOphth, NIHR Biomedical Research Centre, James E. Morgan, DPhil, Cardiff University,
445 Zaynah A. Muthy, BSc, NIHR Biomedical Research Centre, Eoin O'Sullivan, MD, King's College
446 Hospital NHS Foundation Trust, Chris G. Owen, PhD, University of London, Praveen Patel, FRCOphth,
447 NIHR Biomedical Research Centre, Euan Paterson, BSc, Queens University Belfast, Tunde Peto, PhD,
448 Queen's University Belfast, Axel Petzold, PhD, UCL, Jugnoo S. Rahi, PhD, UCL Great Ormond Street
449 Institute of Child Health, Alicja R. Rudnik, PhD, University of London, Jay Self, PhD, University of
450 Southampton, Sobha Sivaprasad, FRCOphth, NIHR Biomedical Research Centre, David Steel,
451 FRCOphth, Newcastle University, Irene Stratton, MSc, Gloucestershire Hospitals NHS Foundation Trust,
452 Nicholas Strouthidis, PhD, NIHR Biomedical Research Centre, Cathie Sudlow, DPhil, University of
453 Edinburgh, Dhanes Thomas, FRCOphth, NIHR Biomedical Research Centre, Emanuele Trucco, PhD,
454 University of Dundee, Adnan Tufail, FRCOphth, NIHR Biomedical Research Centre, Veronique Vitart,
455 PhD, University of Edinburgh, Stephen A. Vernon, DM, Nottingham University Hospitals NHS Trust,

456 Ananth C. Viswanathan, FRCOphth, NIHR Biomedical Research Centre, Cathy Williams, PhD,
457 University of Bristol, Katie Williams, PhD, King's College London, Jayne V. Woodside, MRCOphth,
458 PhD, Queen's University Belfast, Max M. Yates, PhD, University of East Anglia, Jennifer Yip, PhD,
459 University of Cambridge, and Yalin Zheng, PhD, University of Liverpool.

460